

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
2 October 2003 (02.10.2003)

PCT

(10) International Publication Number  
WO 03/080591 A1(51) International Patent Classification<sup>7</sup>: C07D 309/10

(21) International Application Number: PCT/SI03/00009

(22) International Filing Date: 17 March 2003 (17.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
P-200200086 26 March 2002 (26.03.2002) SI(71) Applicant (for all designated States except US): KRKA  
TOVARNA ZDRAVIL, D.D., NOVO MESTO [SI/SI];  
Smarjeska cesta 6, 8501 Novo mesto (SI).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


## Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(74) Agent: PATENTNA PISARNA D.O.O.; Copova 14, POB 1725, 1001 Ljubljana (SI).

(54) Title: PROCESS FOR THE PREPARATION OF 4-OXYTETRAHYDROPYRAN-2-ONES



(I)

(57) Abstract: A process for the preparation of inhibitors of HMG-CoA reductase, such as simvastatin, from 4-silyloxytetrahydropyran-2-ones with triethylamine trihydrofluoride being used as the desilylation reagent is described. The reaction is performed in organic solvents, a mixture thereof or without solvents. It is characteristic of this reaction that no additional impurities are obtained and that it takes place without the use of additional catalysts and with low excesses of the reagent.

WO 03/080591 A1

BEST AVAILABLE COPY